



TITLE:

<ORIGINAL REPORTS>A STUDY ON CHARACTERISTICS OF ATYPICAL MYCOBACTERIA ISOLATED FROM SPUTUM SPECIMENS IN KOREA

AUTHOR(S):

LEE, JayQ.; KIM, SungChin

CITATION:

LEE, JayQ. ...[et al]. <ORIGINAL REPORTS>A STUDY ON CHARACTERISTICS OF ATYPICAL MYCOBACTERIA ISOLATED FROM SPUTUM SPECIMENS IN KOREA. 京都大学結核胸部疾患研究所紀要 1974, 7(2): 98-105

ISSUE DATE:

1974-03-30

URL:

<http://hdl.handle.net/2433/52258>

RIGHT:

ORIGINAL REPORTS

A STUDY ON CHARACTERISTICS OF ATYPICAL MYCOBACTERIA ISOLATED FROM SPUTUM SPECIMENS IN KOREA

Jay Q. LEE and Sung-Chin KIM,

Seoul, Korea

(Received for publication on December 24, 1973)

INTRODUCTION

In many countries of the world, studies on atypical acid-fast bacilli and their bacteriological and clinical significance have become an important field of research since Buhler and Pollak first reported human infection with "atypical acid-fast organisms" in 1953.¹⁾

Interest in this field in Korea was initiated in 1960 when Lee reported isolation of 28 strains of atypical acid-fast bacilli from the sputum cultures of the patients with "active pulmonary tuberculosis".²⁾

In 1970, Kim et al. reported isolation of 97 strains of atypical mycobacteria from more than 22,000 sputum samples from Seoul and its vicinity,³⁾ but, unfortunately, there was hardly any mention on the virulence or clinical significance of these organisms in any of these reports in Korea.

Since the pathogenicity of atypical mycobacteria isolated in Korea is in doubt, the author carried out identification procedures and experiments on the virulence of atypical mycobacteria for the white inbred mice.

MATERIALS AND METHODS

Fifteen strains of atypical mycobacteria were obtained from the acid-fast bacilli cultures in the Central Laboratory of the Korean Tuberculosis Association from February through May, 1970. These strains were isolated on 3% Ogawa medium from sputa following digestion with 4% NaOH or on Löwenstein-Jensen medium following digestion with oxalic acid, and only those strains yielding more than 20 colonies were used in the experiment.

These strains were classified according to Runyon's classification into photochromogens (Group I), scotochromogens (Group II), nonchromogens (Group III) and rapid growers (Group IV).⁴⁾ Subtyping of the organisms were done on the bases of the growth rate, pigmentation of

the colonies, colouring response upon exposure to light ^{5,6)}, niacin test ^{7,8)}, nitrate reduction ⁹⁾, catalase activity ¹⁰⁾, Tween 80 hydrolysis ¹¹⁾, and arylsulfatase activity ¹²⁾. Identification of the organisms was done on the basis of several methods of reference. ^{13,14,15)}

The reference strains utilized in the identification were human type (H37Rv TMC # 102), bovine type (TMC # 401, Revanel), avian type (TMC # 706 McKee # 1) and other atypical mycobacteria, *M. Kanasii* (TMC # 1201, Forbes, p-1), and *M. fortuitum* (TMC # 1530, Martin, PHS # 383), obtained from the mycobacterial culture collection of the Trudeau Institute of the United States.

The experimental animals used were male mice of SM strain (body weight, 13 ± 1 g), and 8 to 10 mice were used for each strain of atypical mycobacteria tested. The inoculum was cultured in Dubos liquid medium for 7 days, and was diluted to 1 mg/ml with distilled water, and 0.2 ml of the latter (average number of viable units, 21.7×10^5) was injected into the tail vein of each mouse.

The following observations were made:

1. Death rate of the animals up to 9 weeks following inoculation.
2. Inspection (*naked eye*) of the pathological changes in the animals sacrificed two at a time from each group at three week intervals up to 9 weeks following inoculation.
3. Weights of the lung and the spleen of the animals thus sacrificed were recorded, and the relative lung weight and the relative spleen weight were calculated according to the following formulae.

$$\text{Relative lung weight} = \frac{\text{lung weight (mg)}}{\text{body weight (g)}} \times 10$$

$$\text{Relative spleen weight} = \frac{\text{spleen weight (mg)}}{\text{body weight (g)}} \times 10$$

4. Weekly colony counts; the entire lungs of the sacrificed mice were put in 2% NaOH, and liquified in universal homogenizer at 15,000 rpm, and 0.1 ml of the material thus obtained was inoculated on each of 10 Löwenstein-Jensen media and was incubated at 37°C.

RESULTS

1. Identification and Classification.

By naked-eye inspection of the isolated colonies, the growth rate and the color were determined, and according to the photochromogen test, the isolated strains were classified according to Runyon's scheme as shown in Table 1. Nine strains (60%) were identified as nonchromogens, 4 were scotochromogens (26.7%) and 2 were rapid growers (13.3%). There were no photo-

Table. 1 Atypical Strains Classified by Runyon's Scheme

| | No. of Strains Isolated | G-I | G-II | G-III | G-IV |
|----------------|-------------------------|-----|------|-------|------|
| No. of Strains | 15 | 0 | 4 | 9 | 2 |
| % | 100.0 | 0 | 26.7 | 60.0 | 13.3 |

chromogens.

The results of various chemical tests were shown in Table 2 to 5. Among the 4 strains of scotochromogenic group, 3 were *M. scrofulaceum*, and 1 was identified as *M. flavescens* showing positive reactions to nitrate reduction test and Tween 80 hydrolysis test. Among the nine of nonchromogenic group, as shown in Table 3, 5 were Battey-avian, 3 were *M. terrae* complex showing positive reaction to nitrate reduction test and Tween 80 hydrolysis test, and 1 was *M. triviale* showing positive reaction to arylsulfatase test. The 2 rapid growers were identified as *M. fortuitum* growing on Löwenstein-Jensen medium within three days and showing positive

Table. 2 Scotochromogenic Group Identified by Various Biochemical Tests

| Strain No. | Growth Within 7 days | Pigment | | Niacin | Nitrate Reduct. | Catalase S.Q. | 68°C PH7 | Tween 80 Hydrolysis | | Arylsulf. | | Species Identified as |
|------------|----------------------|---------|---|--------|-----------------|---------------|----------|---------------------|------|-----------|-----|------------------------|
| | | D. | L | | | | | 5dy | 10dy | 3dy | 2wk | |
| 443 | — | Y | Y | — | — | 50 | + | — | — | — | — | <i>M. scrofulaceum</i> |
| 288 | — | Y | Y | — | — | 50 | + | — | — | — | — | " |
| 183 | — | Y | Y | — | — | 50 | + | — | — | — | ± | " |
| 535 | ± | Y | Y | — | +++ | 50 | + | ± | + | — | ± | <i>M. flavescens</i> |

Note : — Negative
 + Trace but very weak positive
 ± Trace but weak positive
 + Positive
 Y Yellow
 D in dark
 L in light
 SQ Semiquantitative

Table. 3 Non-Chromogenic Group Identified by Various Biochemical Tests

| Strain No. | Growth Within 7 days | Pigment | | Niacin | Nitrate Reduct. | Catalase S.Q. | 68°C PH7 | Tween 80 Hydrolysis | | Arylsulf. | | Species Identified as |
|------------|----------------------|---------|---|--------|-----------------|---------------|----------|---------------------|------|-----------|-----|--------------------------|
| | | D | L | | | | | 5dy | 10dy | 3dy | 2wk | |
| 758 | — | — | — | — | — | 40 | + | — | — | — | — | Battey-avian |
| 447 | — | — | — | — | — | 40 | + | — | — | — | — | " |
| 620 | — | — | — | — | — | 40 | + | — | ± | — | — | " |
| 274 | — | — | — | — | ± | 40 | + | — | — | — | — | " |
| 333 | — | — | — | — | — | 40 | + | — | — | — | ± | " |
| 1278 | — | — | — | — | +++ | 40 | + | ± | + | — | — | <i>M. terrae</i> Complex |
| 906 | — | — | — | — | +++ | 40 | + | ± | + | — | — | " |
| 61 | — | — | — | — | +++ | 40 | + | + | + | — | — | " |
| 130 | — | — | — | — | +++ | 40 | + | ± | + | ± | + | <i>M. triviale</i> |

Table. 4 Rapid Growers Group Identified by Various Biochemical Tests

| Strain No. | Growth Within 7 days | Pigment | | Niacin | Nitrate Reduct. | Catalase S.Q. | 68°C PH7 | Tween 80 Hydrolysis | | Arylsulf. | | Species Identified as |
|------------|----------------------|---------|---|--------|-----------------|---------------|----------|---------------------|------|-----------|-----|-----------------------|
| | | D | L | | | | | 5dy | 10dy | 3dy | 2wk | |
| 12 | + | — | — | — | +++ | 50 | + | — | ± | + | + | <i>M. fortuitum</i> |
| 159 | + | — | — | — | +++ | 50 | + | — | — | + | + | " |

Table. 5 Result of Observations on Mice Infected with Atypical Mycobacteria

| Species | Strain No. | No. of Mice Observed | Viable Unit Infected ($\times 10^5$) | Death Rate | Macroscopic Findings | L/B Weight Ratio | S/B Weight Ratio |
|-----------------------|------------|----------------------|--|------------|----------------------|------------------|------------------|
| <i>M. scrofla.</i> | 443 | 10 | 48 | 0 | 0 | 122 | 135 |
| " | 288 | 10 | 38 | 0 | 0 | 110 | 89 |
| " | 183 | 10 | 19 | 0 | 0 | 129 | 118 |
| <i>M. flaves.</i> | 545 | 9 | 24 | 0 | 0 | 120 | 103 |
| Batt-avium | 759 | 10 | 28 | 0 | 0 | 139 | 65 |
| " | 447 | 10 | 21 | 0 | 0 | 140 | 83 |
| " | 620 | 10 | 27 | 0 | 0 | 104 | 112 |
| " | 274 | 8 | 18 | 0 | 0 | 134 | 87 |
| " | 333 | 10 | 33 | 0 | 0 | 108 | 89 |
| <i>M. terrae</i> | 1278 | 10 | 18 | 0 | 0 | 98 | 65 |
| " | 906 | 10 | 19 | 0 | 0 | 118 | 66 |
| " | 61 | 10 | 11 | 0 | 0 | 130 | 59 |
| <i>M. triviale</i> | 130 | 10 | 6 | 0 | 0 | 127 | 123 |
| <i>M. fortuit</i> | 12 | 10 | 14 | 0 | 0 | 112 | 112 |
| " | 159 | 10 | 22 | 0 | 0 | 88 | 70 |
| <i>M. bovis</i> * | | 10 | 13 | 100. | ++ | 718 | 162 |
| <i>M. avium</i> * | | 10 | 18 | 0 | 0 | 86 | not tested |
| <i>M. kansasii</i> * | | 10 | 10 | 0 | 0 | 103 | " |
| <i>M. fortuit.</i> * | | 10 | 36 | 0 | 0 | 112 | 54 |
| <i>M. TB(H37Rv)</i> * | | 10 | 18 | 90 | ++ | 620 | 141 |
| Control | | 8 | 0 | 0 | 0 | 116 | 44 |

Note : *Reference Strain
Control not infected

L/B Lung and Body
S/B Spleen and Body

Table. 6 Viable Units of Atypical Mycobacteria in Experimental Mice Lungs at Three-Week Intervals

| Species | Strain No. | After 24 hr | 3rd wk | 6th wk | 9th wk |
|--------------------|------------|-------------|--------|--------|--------|
| <i>M. scrofla.</i> | 443 | 56,200 | 0 | 0 | 0 |
| " | 288 | 34,000 | 0 | 0 | 0 |
| " | 183 | 14,200 | 0 | 0 | 0 |
| <i>M. flaves.</i> | 545 | 8,600 | 0 | 0 | 0 |
| Batt-avium | 759 | 50 | 0 | 0 | 0 |
| " | 447 | 34,000 | 1,500 | 0 | 0 |
| " | 620 | 35,500 | 160 | 0 | 0 |
| " | 274 | 68,500 | 1,100 | 220 | 86 |
| " | 333 | 42,500 | 1,000 | 21 | 0 |
| <i>M. terrae</i> | 1,278 | 78,500 | 12,500 | 3 | 0 |
| " | 906 | 12,700 | 6,300 | 11 | 1 |
| " | 61 | 92,700 | 230 | 0 | 0 |
| <i>M. triviale</i> | 130 | 97,500 | 1,500 | 0 | 0 |
| <i>M. fortuit.</i> | 12 | 32,500 | 8 | 0 | 0 |
| " | 159 | 600 | 0 | 0 | 0 |

reaction to aroylsulfatase test.

2. Pathogenicity for mice.

Table 5 shows the results of 9 weeks observation of the mice inoculated with the atypical acid-fast bacteria and the reference strains. None of the experimental animals died within the 9 weeks of observation except those inoculated with the reference strains of H37Rv and *M. bovis*. Also the organs of the sacrificed mice did not show any pathological changes except those from the animals infected with the reference strains of H37Rv and *M. bovis*, showing severe granulomatous changes in the lungs. There was no appreciable difference in the relative lung weights between the study and the control groups, but only the spleen weight of the study group increased as compared with the uninfected control group.

In order to observe the persistence of the bacteria in the lungs, the viable units were counted at 24 hours, 3 weeks, 6 weeks, and 9 weeks as shown in Table 6, and there was a rapid decrease in the numbers of bacteria with the passage of time. *M. scrofulaceum* and *M. flavus* completely disappeared from the lungs within 3 weeks, but other groups of organisms showed persistence at 3rd week in considerable numbers. At 6th week, only a few colonies of Battey-avium 274, 333, and *M. terrae* 1,278 and 906 were obtained, but other strains completely disappeared, and at 9th week only Battey-avium 274, and *M. terrae* 906 survived in small numbers whereas all other strains disappeared from the lung tissue.

DISCUSSION

Since the first report of isolation of atypical mycobacteria from the sputum cultures of suspected tuberculosis patients in Korea in 1960,²⁾ epidemiological and bacteriological works have been published in this country. They uniformly reported recovery of many strains of atypical acid-fast bacilli belonging to Groups II, III, and IV according to Runyon's classification^{2,3)}

However, little information is available as to the clinical significance and virulence of the isolated atypical mycobacteria in Korea. The author's work, up to the time of this writing, represents the only animal experiment to evaluate the virulence of the atypical mycobacteria isolated from human sputa in Korea.

The white mice were used in this study for the possible superiority of the mice to the guinea pigs as experimental animals for the virulence study of atypical mycobacteria.⁴⁾

The isolated strains of atypical mycobacteria in this study belonged to Group II, (26.7%), Group III (60%) and Group IV (13.3%) according to Runyon's classification. This is generally similar to the findings of other workers in Korea, especially the notable absence of the strains belonging to Group I, which are considered to be more virulent than others,^{2,3)} and the relative preponderance of strains belonging to Group III.^{3,16)}

As shown in Table 5, none of the strains of atypical mycobacteria tested caused death of the infected mice within the nine week period of observation, and only the relative spleen weight somewhat increased in the study group. The number of viable units cultured from the lung tissue of the infected mice rapidly decreased with the passage of time, and almost completely disappeared by the 9th week of observation.

In short, the fifteen strains of atypical mycobacteria belonging to Groups II, III and IV recovered from the sputum cultures of the radiologically diagnosed cases of "pulmonary tuberculosis" failed to cause demonstrable disease in mice when inoculated intravenously. The results of this study could at least partly be explained by the fact that there was no strain ascribable to Group I (photochromogen) which is generally considered to be the most virulent of all the atypical mycobacteria.⁴⁾ It is interesting to note, in this connection, that there is some similarity in the group distribution between the strains of atypical mycobacteria isolated from the soil samples¹⁶⁾ and those obtained from the human sputum cultures in Korea,³⁾ which show relative preponderance of Group III strains and absence of Group I strains. It is also noted, in this context, that atypical mycobacteria were cultivated from sputum samples of 7.5% of 290 apparently healthy men in tropical Australia, and most of the strains belonged to Groups II and III.¹⁷⁾ However, the possibility remains that these apparently innocuous organisms may cause opportune infection depending upon the condition of the host.

Another point to consider is that the author's experimental model including the route of inoculation may be unsatisfactory. While the author's experimental animals did not show any sign of disease, Blitek-Golc reported 50% mortality in the mice infected with Group III strains, but the route of inoculation was not mentioned in the abstract.¹⁸⁾ Japanese workers reported successful production of tuberculous lesions in the testicles of rabbits inoculated with atypical mycobacteria,¹⁹⁾ and even in guinea pigs inoculated by intraepididymal route.²⁰⁾ Therefore, it is possible that pathological lesions could have been produced in the author's mice if various other routes of inoculation had been utilized.

Still another point to consider is finding the most suitable experimental animal for virulence tests of atypical mycobacteria. The success of any experimental animal inoculation of mycobacteria depends to a considerable extent upon the choice of an appropriate test animal. Recently, Choi et al. showed that Korean chipmunk proved to be a reliable experimental animal for *M. tuberculosis* even with the lower infectious dose (less than 5.8×10^3), the establishment of infection becoming manifest in the second week of inoculation.²¹⁾ In the author's opinion, further studies with atypical mycobacteria should be tried not only by various routes of inoculation in mice but also on some other experimental animals such as Korean chipmunk.

SUMMARY AND CONCLUSION

Fifteen strains of atypical mycobacteria obtained from sputum cultures of clinical cases of "tuberculosis" were classified into Groups II, III and IV by Runyon's classification. There was notable absence of Group I strains.

None of the test strains inoculated into mice by intravenous route for virulence test produced pathological lesions in mice, and the atypical mycobacteria rapidly disappeared from the lung tissue of the animals.

It is concluded that these strains of atypical mycobacteria from the sputum samples in Korea were of no significant pathogenicity in mice when inoculated by the tail vein, and that future trials should utilize other routes of inoculation or other experimental animals such as Korean chipmunk.

ACKNOWLEDGEMENT

The authors wish to express their gratitude to professor I. Uesaka of Kyoto University for his scholarly advices in the course of this study and his critical review of the manuscript.

REFERENCES

- 1) Buhler, V. B., and Pollak, A.: Human infection with atypical acid-fast organisms: Report of two cases with pathological findings. *Amer. J. Clin. Path.*, 23: 363, 1953.
- 2) Lee, W. K.: The biological study of atypical acid-fast bacilli. *Tuberculosis (Korea)*, 7: 66, 1960.
- 3) Kim, S. C., and Kim, S. C.: A study on unclassified mycobacteria isolated from human sputa. *Tuberc. Resp. Dis. (Korea)*, 17: 33, 1970.
- 4) Runyon, E. H.: Anonymous mycobacteria in pulmonary disease. *Med. Clin. North Amer.*, 43: 273, 1959.
- 5) Fregnan, G. C. and Smith, D. W.: Description of various colony forms of mycobacteria. *J. Bact.*, 83: 819, 1962.
- 6) Vestal, A. L., and Kubica, G. P.: Differential colonial characteristics of mycobacteria on Middlebrook and Cohen 7H-10 agar base medium. *Am. Rev. Resp. Dis.*, 94: 247, 1966.
- 7) Konno K.: New chemical method to differentiate human-type tubercle bacilli from other mycobacteria. *Science*, 124: 985, 1956.
- 8) Runyon, E. H., Selin, M. J., and Harris, H. W.: Distinguishing mycobacteria by the niacin test. *Am. Rev. Resp. Dis.*, 79: 663, 1959.
- 9) Virtanen, S.: A study of nitrate reduction by mycobacteria. *Acta Tuberc. Scand.*, suppl. 48, 1960.
- 10) Kubica, G. P., et al.: Differential identification of mycobacteria. I. Test on catalase activity. *Am. Rev. Tuberc.*, 94: 400, 1966.
- 11) Wayne, L. G., Doubek, J. R. and Russell, R. L.: Classification and identification of mycobacteria. I. Test employing Tween 80 as substrate. *Am. Rev. Resp. Dis.*, 90: 588, 1964.
- 12) Whitehead, J. E. M., Wildy, R. and Engback, H. C.: Arylsulfatase activity of mycobacteria. *J. Path. Bact.*, 65: 451, 1953.
- 13) Jones, W. D., Abbot, V. D., and Vestal, A. L., Kubica, G. P.: A hitherto undescribed group of non-chromogenic mycobacteria. *Am. Rev. Resp. Dis.*, 94: 790, 1965.
- 14) Sommers, H. M. and Russell, J. P.: The clinically significant mycobacteria: Their recognition and identification. *J. A. S. C. P.*, 1967.
- 15) Vestal, A. L.: Procedures for the isolation and identification of mycobacteria. *Nat'l. Center for Disease Control*, 1969.
- 16) Kim, S. C. et al.: A study on the acid-fast bacteria isolated from soil in Korea. *Tuberc. Resp. Dis.*, 18: 19, 1971.
- 17) Kiewiet, A. A., and Thompson, J. E.: Isolation of 'atypical' mycobacteria from healthy individuals in tropical Australia. *Tubercle*, 51: 296, 1970.
- 18) Blitek-Golc: Studies on the virulence of atypical mycobacteria isolated from inpatients of the clinic. *Gruzlica*, 37: 409, 1969, cited from abstracts, *Am Rev. Resp. Dis.*, 101: 167, 1970.
- 19) Nakano, H.: Inoculation of atypical mycobacteria into testicles of rabbits. *Kekkaku*, 45: 61, 1970.
- 20) Sushida, K. et al.: Pathogenicity of unclassified mycobacteria for guinea pigs. *Kekkaku*, 45: 339, 1970.
- 21) Choi, T. K. et al.: Determination of minimum infectious dose of *M. tuberculosis* in experimental tuberculosis on Korean chipmunk. *Yonsei J. Med. Sci.*, 5: 1, 1972.

韓国において喀痰から分離された 非定型抗酸性菌の特長について

(大韓民国ソウル市)

李 在 珪, 金 成 鎮

韓国で1970年2月より5月迄に喀痰から分離された抗酸性菌の内、15株の非定型抗酸性菌の性状を検討した。その結果、Runyon 分類で第1群に属する菌株はなく、第2群4株3株 (*M. scrofulaceum* 3株, *M. flavescentis* 1株), 第3群9株 (Battey-avium 型5株, *M. terrae* 群3株, *M. triviale* 1株), 及び第4群2株 (*M. fortuitum*) が認められた。第1群がなく、第3

群が最も多い点は従来韓国の報告と一致する。

更に上記15株につき、その $10^5 \sim 10^6$ をマウス静脈に接種したが、9週間の間に死亡マウスはなく、肺内の生菌数も速かに減少した。非定型抗酸性菌の毒力検定には上記の如き慣用法以外の方法、例えば動物を変える (Korean chipmunk を使用する) 等をとるべきではないかと考える。